



BOKSMART 2010

**ON-FIELD AIRWAY MANAGEMENT
TECHNIQUES**

Providing coaches, referees, players, and administrators with the knowledge, skills, and leadership abilities to ensure that safety and best practice principles are incorporated into all aspects of contact rugby.



WWW.BOKSMART.COM



A joint initiative by SARU and the Chris Burger/Petro Jackson Fund

AIRWAY MANAGEMENT WITH POSSIBLE CERVICAL INJURY
HEAD TILT CHIN LIFT VERSUS CHIN LIFT

Prof. Robert Dunn

MBChB (UCT) MMed (UCT) Orth, FCS(SA) Orth

Consultant Spine and Orthopaedic Surgeon

Head: Orthopaedic Spine Services – Groote Schuur Hospital, Cape Town

Clinical Adviser to BokSmart programme

SUMMARY:

The 'chin lift' or 'jaw thrust' technique should be used for on-field airway control. Any head tilt should be avoided.

INTRODUCTION:

Rugby is a contact sport and players are therefore at risk of injury. A player sustaining a head injury may experience a reduced level of consciousness. At this time the tongue may fall back and obstruct breathing. Primary medical care personnel are trained to protect the airway as priority. Failure to do so may result in asphyxiation and death.

In a collision there is a possibility of cervical spine injury. It is frequently associated with head injury and transient reduced level of consciousness. Thus although airway management is a priority, it should be performed in a manner that minimises cervical motion and thus avoids secondary spinal cord injury in a potentially unstable spine.

DEFINITION AND MECHANISM:

CHIN LIFT'

This manoeuvre lifts the chin forward without changing neck or head position. Fingers of one hand are placed under the mandible, which is gently lifted to bring the chin forward. The thumb of the same hand simultaneously depresses the lower lip to open the mouth. If necessary, the thumb may be placed behind the lower teeth and the chin lifted gently. The neck should never be hyperextended.

JAW THRUST'

The angles of the jaw are grasped on both sides, with each hand as well as the mandible brought forward. This requires both hands and is useful when administering a ventilation mask.

HEAD TILT

This is when the head is hyperextended and is contra-indicated in any patient with a potential cervical injury. This is generally used when intubating a patient (with a stable neck) to allow access to the trachea.

LITERATURE REVIEW:

Manipulation of the head has an effect on the cervical spine. Mechanistic studies have been done to quantify this. Donaldson² investigated the effect of destabilising the C5/6 vertebral level in a cadaver model. They assessed that the intact spine moved an average of 8.76° at C5/6 during flexion and extension. Once the level was destabilised, this increased to 45.5°. Likewise, the translation increased from 0.43mm to 14.18mm. This confirms the intuitive advice to minimise cervical motion in the setting of instability, as these exaggerated movements will increase the forces applied to the spinal cord. They went further to assess various airway manipulation and intubation techniques. Chin lift and jaw thrust still caused some motion at the unstable level, viz. 3.51° and 0.57mm. These are close to the motion tolerated by the normal spine.

Brimacombe³ performed a similar study in cadavers, destabilising the C3 level. They recorded a maximum displacement of 1.9mm and angular change of 2.7° with the chin lift / jaw thrust technique.

Meier⁴ studied the effect of various airway maintenance techniques in anaesthetised children. She found the chin lift and jaw thrust to be similarly effective. Guildner⁵ however reported that the chin lift technique provided the most consistently adequate airway when assessing anaesthetised elective paediatric patients. Roth assessed the head-tilt versus jaw-lift techniques in the paediatric group while anaesthetised but breathing spontaneously. He found the head tilt to be ineffective in 12% and recommended the jaw thrust⁶.

Swartz⁷ et al published the National Athletic Trainers' Association Position Statement (USA) in 2009 after an extensive literature review. This comprehensive document deals with many aspects of care. As regards airway management they are quite clear on promoting the jaw thrust manoeuvre over the head-tilt technique, as the latter produces unnecessary motion at the head and in the cervical spine. They allocated this a "B" category of evidence, which meant that their recommendation was based on inconsistent or limited-quality patient-oriented evidence.

PROS AND CONS:

It is clear both intuitively and from the available literature that airway management should limit cervical motion as much as possible in the known or suspected cervical spine injury.

When a collision occurs on the rugby field, a cervical spine injury must be considered until it can be excluded. Should the player complain of any neck pain, s/he should be managed as if a cervical injury has occurred.


In the case of concomitant head injury and reduced level of consciousness, the neck must again be managed as if an injury exists.

It is quite clear that the 'head tilt' technique causes cervical motion and confers a risk of spinal cord injury should a cervical injury be present. In contrast, cadaver studies have confirmed minimal cervical motion with the chin lift / jaw thrust techniques.

Thus the airway should be maintained by the chin lift or jaw thrust technique.

In the unlikely event that this fails to maintain the airway, and only then, gentle head tilt may be used based on the philosophy of "life over limb".

HEAD TILT – CHIN LIFT TECHNIQUE (not recommended as cervical spine at risk!)



CHIN LIFT TECHNIQUE



JAW THRUST TECHNIQUE



REFERENCES:

- 1) ATLS Student course manual 7th Edition. American College of Surgeons. 2004
- 2) Donaldson WF, Towers JD, Doctor A et al. A Methodology to evaluate motion of the unstable spine during intubation techniques. Spine 1993 18(14):2020-23
- 3) Brimacombe J, Keller C, Ku K et al. Cervical Spine Motion During Airway Management: A Cinefluoroscopic Study of the Posteriorly Destabilized Third Cervical Vertebrae in Human Cadavers. Anesth Analg 2000;91:1274–8
- 4) Meier S, Geiduschek J, Paganoni R et al. The Effect of Chin Lift, Jaw Thrust, and Continuous Positive Airway Pressure on the Size of the Glottic Opening and on Stridor Score in Anesthetized, Spontaneously Breathing Children. Anesth Analg 2002;94:494–9
- 5) Guildner C. Resuscitation — Opening the airway: A comparative study of techniques for opening an airway obstructed by the tongue. Journal of the American College of Emergency Physicians 1976; 5(8): 588-590
- 6) Roth B, Magnusson J, Johansson I et al. Jaw lift — a simple and effective method to open the airway in children. Resuscitation 39 (1998) 171–174
- 7) Swartz E, Boden B, Courson et al. National Athletic Trainers' Association Position Statement: Acute Management of the Cervical Spine–Injured Athlete. Journal of Athletic Training 2009;44(3):306–331

